



## **Polymorphism in the fatty acid desaturase genes and diet are important determinants of infant n-3 fatty acid status.**

**Harsløf, L.B.S.; Larsen, L.H.; Ritz, C.; Hellgren, Lars; Michaelsen, K. F.; Vogel, U.; Lauritzen, L.**

*Publication date:*  
2013

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

### *Citation (APA):*

Harsløf, L. B. S., Larsen, L. H., Ritz, C., Hellgren, L., Michaelsen, K. F., Vogel, U., & Lauritzen, L. (2013). *Polymorphism in the fatty acid desaturase genes and diet are important determinants of infant n-3 fatty acid status..* Abstract from IUNS 20th International Congress of Nutrition, Granada, Spain.

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# **POLYMORPHISM IN THE FATTY ACID DESATURASE GENES AND DIET ARE IMPORTANT DETERMINANTS OF INFANT N-3 FATTY ACID STATUS.**

Harsløf LBS<sup>1</sup>, Larsen LH<sup>1</sup>, Ritz C<sup>1</sup>, Hellgren LI<sup>2</sup>, Michaelsen KF<sup>1</sup>, Vogel U<sup>3</sup> & Lauritzen L<sup>1</sup>

<sup>1</sup>Department of Nutrition, Exercise & Sports, Faculty of Science, University of Copenhagen, Rolighedsvej 30, 1958 Frederiksberg C, Denmark. Telephone: +45 3533 2508, Fax: +45 3533 2483 and E-mail: ll@life.ku.dk

<sup>2</sup>Center for Biological Sequence Analysis, Department of Systems Biology, Technical University of Denmark, Denmark

<sup>3</sup>National Research Centre for the Working Environment, Denmark

**Background and objectives:** Tissue docosahexaenoic acid (DHA) accretion in early infancy has been shown to be supported by the DHA-content of breast-milk and thus may decrease once complementary feeding takes over. Endogenous synthesis of DHA from alpha-linolenic acid has been shown to be very low and polymorphism in the genes that encodes the fatty acid desaturases (FADS) has little effect on DHA-status in adults. It is however unclear to what extent endogenous DHA-synthesis contributes to infant DHA-status.

**Aim:** To investigate the role of diet and FADS polymorphism on DHA-status at 9 months and 3 years.

**Methods:** This cross-sectional study with Danish infants use data from two prospective studies (EFiON and the SKOT cohort). We measured erythrocyte (RBC) DHA-status at 9 months (n=409) and 3 years (n=176) and genotyped 4 FADS tagSNPs, rs3834458, rs1535, rs174575 and rs174448 (n=401).

Information about breastfeeding was obtained by questionnaires and fish intake was assessed by 7-day pre-coded food diaries.

**Results:** FADS-genotype, breastfeeding, and fish intake were found to explain 25% of the variation in infant RBC DHA-status (mean±SD: 6.6±1.9% of the fatty acids (FA%)). Breastfeeding was the most important contributor and still being breast-fed at 9 months was associated with 0.8 FA% higher DHA vs. no longer breast-fed (p<0.001). Two of the examined FADS-SNPs were highly correlated (rs1535 and rs3834458; r=0.98). Homozygous carriers of the minor allele of rs1535 had an increase in RBC DHA of 1.6 FA% relative to those with wild type, whereas minor allele carriers of rs174448 and rs174575 had a decrease of 0.9 (p=0.017) and 1.9 FA% (p=0.001), respectively. Each 10-gram increment in fish intake was associated with an increase in DHA-status of 0.3 FA%. At 3 years, fish intake was the only significant determinant of DHA-status (0.2 FA%/10g).

**Conclusions:** FADS-genotype and diet are both important determinants of DHA-status in late infancy.

**Key words:** FADS-genotype, docosahexaenoic acid, fish intake, child nutrition, breastfeeding